

**CORRECTIONS TO**  
*A RADICAL APPROACH TO LEBESGUE'S THEORY OF INTEGRATION*

**Page 10**, line 4, “Propogation” should be “Propagation”

**Page 12**, Footnote 5, “Weierstrass’s” should be “Weierstrass.”

**Page 23**, line before displayed quote, it was to Riemann’s father, rather than to Richard Dedekind, to which this letter was addressed

**Page 23**, footnote, the Hochkirchen reference should be to page 263 rather than 261.

**Page 24**, line 8, “oscillirendend” should be “oszillierenden.”

**Page 27**, Theorem 2.1, line 4, change “We can find a  $\delta$  response” to “We can find a response  $\delta$ ”

**Page 30**, last set of displayed equations, second last line should be

$$= \lim_{\epsilon_1 \rightarrow 0^-} \left( -2|\epsilon_1|^{1/2} + 2 \right) + \lim_{\epsilon_2 \rightarrow 0^+} \left( 2 - 2|\epsilon_2|^{1/2} \right)$$

**Page 38**, first 6 lines (the justification that  $w(x)$  is not differentiable at  $a_N$ ). This justification is not correct. The correct justification is given by:

We first observe that  $\sum_{n=1}^{N-1} k^n h(x - a_n)$  is differentiable at  $x = a_N$ . Therefore,  $w$  is differentiable at  $x = a_N$  if and only if  $\sum_{n=N}^{\infty} k^n h(x - a_n)$  is differentiable at  $x = a_N$ . If this sum is differentiable, then the oscillation of the average rate of change between  $a_N$  and  $x$ , as a function of  $x$ , can be made arbitrarily close to 0 by taking  $x$  sufficiently close to (but not equal to)  $a_N$ . But this average rate of change can be written as a sum of two fractions:

$$\frac{k^N h(x - a_N) - k^N h(0)}{x - a_N} + \frac{\sum_{n=N+1}^{\infty} k^n h(x - a_n) - \sum_{n=N+1}^{\infty} k^n h(a_N - a_n)}{x - a_N}.$$

The first fraction,

$$\frac{k^N h(x - a_N) - k^N h(0)}{x - a_N} = k^N \left( 1 + \frac{1}{2} \sin \left( \frac{1}{2} \ln [(x - a_N)^2] \right) \right),$$

oscillates between  $k^N/2$  and  $3k^N/2$  over any interval that contains  $a_N$ . By the mean value theorem,

$$\left| \frac{\sum_{n=N+1}^{\infty} k^n h(x - a_n) - \sum_{n=N+1}^{\infty} k^n h(a_N - a_n)}{x - a_N} \right| < 2 \sum_{n=N+1}^{\infty} k^n = \frac{2k^{N+1}}{1 - k},$$

---

*Date:* August 20, 2009.

and therefore the oscillation of the second fraction is strictly less than  $4k^{N+1}/(1-k)$ . If  $k < 1/5$ , then the oscillation of the second fraction is bounded above by an amount strictly less than the oscillation of the first fraction, and therefore the sum of their oscillations cannot be made arbitrarily small.

**Page 39**, line 4, “Exercises 2.2.4–2.2.5” should be “Exercises 2.2.4–2.2.9”

**Page 42**, line 2 under **Hankel’s Innovations**, “Funktionen” should be “Funktionen.”

**Page 46**, paragraph 3, line 6: “funczioni” should be “funzioni”  
 paragraph 4, line 2, “Riechen” should be “Reihen”

**Page 47**, third definition box, “A  $S$  set is type  $n$ ” should be “A set  $S$  is type  $n$ ”

**Page 50**, Exercise 2.3.13,  $\sin(1/f_1(x))$  and  $\sin(1/f_2(x))$  should be  $\sin(1/f_1(x))$  and  $\sin(1/f_2(x))$

**Page 50**, Exercise 2.3.15, “Show that if  $C \in \mathcal{C}$  is a finite cover” should be “Show that if  $C \in \mathcal{C}$  is a finite cover”

**Page 55**, last paragraph of proof of Theorem 3.1, change “then pick any  $f(x) \in T$  and any  $\epsilon$ ” to “then pick any  $x$  and any  $\epsilon$ ”

**Page 68**, third full paragraph, line 3, “Functionenlehre” should be “Funktionenlehre”

**Page 70**, Exercise 3.2.12, the displayed summations

$$\sum_{n=1}^{\infty} \frac{A_n}{b_n} \quad \text{and} \quad \sum_{n=1}^{\infty} b_n$$

should be

$$\sum_{n=1}^{\infty} \frac{A_n}{b_n} \quad \text{and} \quad \sum_{n=1}^{\infty} b_n$$

**Page 81** fifth line from bottom: “there are also nowhere dense set that” should be “there are also nowhere dense sets that”.

**Page 86**, last paragraph, change

This function has the curious property that it is a nonconstant function with a derivative that exists and is zero at every point except at a set with outer content zero.

to

This function has the curious property that it is a continuous nonconstant function with a derivative that exists and is zero at every point except at a set with outer content zero.

**Page 91**, graph of SVC(4): The top row of line segments should each have their middle quarters removed.

**Page 98**, last paragraph, line 6, “Functionenlehre” should be “Funktionenlehre”

**Page 112**, second line from bottom: “Dirichet’s function” should be “Dirichlet’s function”.

**Page 114**, line 1, “The function  $g$  of Exercise 1.1.5 (p. 15)” should be “The function  $m$  of Exercise 2.1.14 (p. 32)”

**Page 126**, third line from top: “well-fined” should be “well-defined”.

**Page 149**, Exercise 5.3.10. Second line. The inequality “ $m_\epsilon(S) < m(U) + \epsilon$ ” should be “ $m(U) < m_\epsilon(S) + \epsilon$ .”

**Page 150**, Exercise 5.3.15, “the supremum and the infimum” should be “the lim sup and the lim inf”

**Page 157**, Exercises 5.4.2 and 5.4.3: For both exercises, we need to assume that the set in question is measurable. Both first sentences should begin, “Show that any measurable set of positive measure contains . . .”

**Page 196**, Proof of Theorem 6.24, the symbol indicating the end of the proof should come just before the next subhead: **Limits of Step Functions**

**Page 203**, End of second paragraph: “functional theorem of calculus” should be “fundamental theorem of calculus.”

**Page 205**, line 2, second set of equalities should be

$$D^- f(0) = D_- f(0) = -1.$$

Paragraph 3, line 3, “partricular” should be “particular”

**Page 208**, line 4 from bottom, “ $f = f - (T - f)$ ” should be “ $f = T - (T - f)$ .”

**Page 220**, Equation (7.16) should be

$$g(t_x) - g(s_x) > N(t_x - s_x).$$

**Page 295**, line 6 from bottom,  $|x_j - x_{j-1}| < \delta(x_j^*) < \inf_{x \in G_k^C} |x_k^* - x|$  should be  $|x_j - x_{j-1}| < \delta(x_j^*) < \inf_{x \in G_{k_j}^C} |x_j^* - x|$ . Also note that the inequalities at the top of page 296 remain valid even when several tags  $x_j^*$  share the same  $E_k$ . It works because each of them generates a separate contribution within the same set  $(G_k - E_k)$ .

**Page 297**, line 18, “most useful” should be “more useful”

**Page 300**, Hint for exercise 1.2.6. First sentence should read, “Let  $S_n = \sup_{k \geq n} a_k$ . ”

**Page 303**, Hint for exercise 3.3.10, line 4. Both occurrences of  $A'$  should just be  $A$ . This line should read: “ $n$  so that  $\psi^n(a) \in B - A$ , then  $a$  is mapped to  $\psi(a)$ . If  $a$  is not in  $A$  or there ...”

**Page 309**., Hint for exercise 6.4.9. The two numbers  $a, b \in \mathcal{N}$  need to be on the same side of  $x$ .

**Page 309**., Hint for exercise 6.4.14. Change the hint to read:

For each  $n \in \mathbb{N}$ , choose a measurable set  $E_n$  with measure less than  $1/n$  for which  $f$  is continuous relative to  $[a, b] - E_n$ . Show that there is an open set  $U_n$  such that

$$U_n - E_n \subseteq \{x \mid f(x) > c\} \subseteq U_n \cup E_n.$$

Use exercise 5.3.9.

**Page 319**., Hochkirchen, T. 2003. Page numbers should be 261–290.

**Thanks to:** Patrice Goyer, Sean Hickey, Peter Horton, Jim Leonard, Stylianos Pezaris, Fred Rickey, Bill Rozzi, Allen Rubenstein, Lou Talman, and Hao Zou.